arXiv:1202.6565v2 [math.CV] 8 Mar 2012

# SHARP LIPSCHITZ CONSTANTS FOR THE DISTANCE RATIO METRIC

SLAVKO SIMIĆ, MATTI VUORINEN, AND GENDI WANG

Abstract. We study expansion/contraction properties of some common classes of mappings of the Euclidean space $\mathbb{R}^n, n \geq 2$, with respect to the distance ratio metric. The first main case is the behavior of Möbius transformations of the unit ball $\mathbb{B}^n$ in $\mathbb{R}^n$ onto itself. In the second main case we study the behavior of bounded analytic functions or polynomials of the unit disk. In both cases sharp constants are obtained.

## 1. Introduction

Conformal invariants and conformally invariant metrics have been some of the key notions of geometric function theory and of quasiconformal mapping theory for several decades. One of the modern trends is to extend this theory to Euclidean spaces of higher dimension or to more general metric spaces and new methods have been invented. It has turned out that we cannot any more expect the same invariance properties as in the classical cases, but still some type of "nearinvariance" or "quasi-invariance" is a desirable feature. The quasihyperbolic metric of a domain $G \subset \mathbb{R}^n, n \geq 2$ is one of these new notions and it is known to be quasi-invariant in the sense described below and there are many reasons why one may regard it as a spatial version of the hyperbolic metric. Another crucial notion, which is the object of the present study, is the distance ratio metric. It is often applied to study quasihyperbolic geometry. Very little is known about the geometric properties of balls in these two spaces: only recently it was proved for instance that in some special cases for small radii the balls are convex [K1, K2, MV, V2, RT] no matter where the center is. This convexity property is expected to hold in general, but it has not been proved yet.

We start with the definitions of some of the metrics relevant for our study.

**Distance ratio metric.** For a subdomain $G \subset \mathbb{R}^n$ and for all $x, y \in G$ the distance-ratio metric $j_G$ is defined as

$$j_G(x,y) = \log\left(1 + \frac{|x-y|}{\min\{d(x,\partial G), d(y,\partial G)\}}\right),$$

where $d(x,\partial G)$ denotes the Euclidean distance from the point $x$ to the boundary $\partial G$ of the domain $G$. The distance ratio metric was introduced by F.W. Gehring and B.P. Palka [GP] and in the above simplified form by M. Vuorinen [Vu1] and it is frequently used in the study of hyperbolic type metrics [HIMPS] and geometric theory of functions. J. Väisälä's theory of quasiconformal maps in infinite dimensional Banach spaces is entirely based on the quasihyperbolic metric [V1].

---

`File: jlipfusion120308.tex, printed: 2012-3-9, 5.13`

2010 *Mathematics Subject Classification.* 51M10(30C20).

*Key words and phrases.* Lipschitz constant, Möbius transformation.

Let $p \in G$, then for all $x, y \in G \setminus \{p\}$

$$j_{G\setminus\{p\}}(x,y) = \log\left(1 + \frac{|x-y|}{\min\{d(x,\partial G), d(y,\partial G), d(x,p), d(y,p)\}}\right).$$

From the above formula it is easy to see that $j$-metric depends on the boundary of the domain highly.

**Absolute ratio metric.** Let $G$ be an open subset of $\mathbb{R}^n$ with card $\partial G \geq 2$. For all $x$, $y \in G$, the absolute ratio metric $\delta_G$ is defined as

$$\delta_G(x,y) = \log(1 + \sup_{a,b\in\partial G} |a,x,b,y|), \quad |a,x,b,y| = \frac{|a-b||x-y|}{|a-x||b-y|}.$$

It is a well-known basic fact that $\delta_{\mathbb{B}^n}$ agrees with the hyperbolic metric both in the case of the unit ball as well as in the case of the half space (cf. [Vu2, 8.39]).

**Quasihyperbolic metric.** Let $G$ be an open subset of $\mathbb{R}^n$. For all $x$, $y \in G$, the quasihyperbolic metric $k_G$ is defined as

$$k_G(x,y) = \inf_\gamma \int_\gamma \frac{1}{d(z,\partial G)}|dz|,$$

where the infimum is taken over all rectifiable arcs $\gamma$ joining $x$ to $y$ in $G$.

The hyperbolic metric in the unit ball or half space, the absolute ratio metric $\delta_G$ are Möbius invariant. However, neither the quasihyperbolic metric $k_G$ nor the distance ratio metric $j_G$ are invariant under Möbius transformations. Therefore, it is natural to ask what the Lipschitz constants are for these metrics under conformal mappings or Möbius transformations in higher dimension. F. W. Gehring, B. P. Palka and B. G. Osgood proved that these metrics are not changed by more than a factor 2 under Möbius transformations, see [GP, Corollary 2.5] and [GO, proof of Theorem 4]:

**Theorem 1.1.** *If $D$ and $D'$ are proper subdomains of $\mathbb{R}^n$ and if $f$ is a Möbius transformation of $D$ onto $D'$, then for all $x, y \in D$*

$$\frac{1}{2} m_D(x,y) \leq m_{D'}(f(x), f(y)) \leq 2 m_D(x,y),$$

*where $m \in \{j, k\}$.*

R. Klén, M. Vuorinen and X. Zhang proposed a conjecture in [KVZ]:

**Conjecture 1.2.** *Let $a \in \mathbb{B}^n$ and $h : \mathbb{B}^n \to \mathbb{B}^n$ be a Möbius transformation with $h(a) = 0$. Then*

$$\sup_{\substack{x,y\in\mathbb{B}^n \\ x\neq y}} \frac{j_{\mathbb{B}^n}(h(x), h(y))}{j_{\mathbb{B}^n}(x,y)} = 1 + |a|.$$

**Remark 1.3.** *The sharpness of the constant is proved in* [KVZ, Remark 3.4] *by taking $x = ta/|a| = -y$, $t \in (0,1)$ and letting $t \to 0^+$.*

In this paper, we find sharp Lipschitz constants for the distance ratio metric under Möbius transformations of the unit ball in $\mathbb{R}^n$ and for $n = 2$ under analytic functions or polynomials of the unit disk.

**Main results**

**Theorem 1.4.** *Let* $a \in \mathbb{B}^n$ *and* $f : \mathbb{B}^n \to \mathbb{B}^n = f(\mathbb{B}^n)$ *be a Möbius mapping with* $f(a) = 0$. *Then for all* $x, y \in \mathbb{B}^n$

$$\frac{1}{1+|a|} j_{\mathbb{B}^n}(x,y) \le j_{\mathbb{B}^n}(f(x), f(y)) \le (1+|a|) j_{\mathbb{B}^n}(x,y),$$

*and the constants* $\frac{1}{1+|a|}$ *and* $1+|a|$ *are both best possible.*

**Theorem 1.5.** *(1) Let* $a \in \mathbb{H}^2$ *and* $f : \mathbb{H}^2 \to \mathbb{B}^2$ *be a Möbius mapping with* $f(a) = 0$. *Then for all* $x, y \in \mathbb{H}^2$

$$\frac{1}{2} j_{\mathbb{H}^2}(x,y) \le j_{\mathbb{B}^2}(f(x), f(y)) \le 2 j_{\mathbb{H}^2}(x,y),$$

*and the constants* $\frac{1}{2}$ *and* $2$ *are both best possible.*

*(2) Let* $a \in \mathbb{H}^2$ *and* $f : \mathbb{H}^2 \setminus \{a\} \to \mathbb{B}^2 \setminus \{0\}$ *be a Möbius mapping with* $f(a) = 0$. *Then for all* $x, y \in \mathbb{H}^2 \setminus \{a\}$

$$\frac{1}{2} j_{\mathbb{H}^2 \setminus \{a\}}(x,y) \le j_{\mathbb{B}^2 \setminus \{0\}}(f(x), f(y)) \le 2 j_{\mathbb{H}^2 \setminus \{a\}}(x,y),$$

*and the constants* $\frac{1}{2}$ *and* $2$ *are both best possible.*

**Theorem 1.6.** *Let* $\{a_k\}$ *be a sequence of complex numbers and* $\sum_{k=0}^{\infty} |a_k| \le 1$. *Let* $f : \mathbb{B}^2 \to \mathbb{B}^2$ *be a non-constant mapping with* $f(z) = \sum_{k=0}^{\infty} a_k z^k$. *Then for all* $x,\ y \in \mathbb{B}^2$

$$j_{\mathbb{B}^2}(f(x), f(y)) \le j_{\mathbb{B}^2}(x,y),$$

*and this inequality is sharp.*

It should be noted that these metrics can be estimated one in terms of the other and the hyperbolic metrics $\rho_{\mathbb{B}^n}$ and $\rho_{\mathbb{H}^n}$ . For instance we have the following inequalities [Vu2, 2.41] and [AVV, 7.56]

$$\frac{1}{2} \rho_{\mathbb{B}^n}(x,y) \le j_{\mathbb{B}^n}(x,y) \le k_{\mathbb{B}^n}(x,y) \le \rho_{\mathbb{B}^n}(x,y) \,, \forall\ x, y \in \mathbb{B}^n \,,$$

$$\frac{1}{2} \rho_{\mathbb{H}^n}(x,y) \le j_{\mathbb{H}^n}(x,y) \le k_{\mathbb{H}^n}(x,y) \equiv \rho_{\mathbb{H}^n}(x,y) \,, \forall\ x, y \in \mathbb{H}^n \,.$$

## 2. Lipschitz Constants under Some Special Mappings

**Möbius transformations.** The group of Möbius transformations in $\overline{\mathbb{R}^n}$ is generated by transformations of two types

(1) reflections in hyperplane $P(a,t) = \{x \in \mathbb{R}^n : x \cdot a = t\}$

$$f_1(x) = x - 2(x \cdot a - t)\frac{a}{|a|^2},\ f_1(\infty) = \infty;$$

(2) inversions (reflections) in $S^{n-1}(a,r) = \{x \in \mathbb{R}^n : |x-a| = r\}$

$$f_2(x) = a + \frac{r^2(x-a)}{|x-a|^2},\ f_2(a) = \infty, f_2(\infty) = a.$$

If $G \subset \overline{\mathbb{R}^n}$ we denote by $\mathcal{GM}(G)$ the group of all Möbius transformations with $fG = G$.

We denote $a^* = \frac{a}{|a|^2}$ for $a \in \mathbb{R}^n \setminus \{0\}$, and $0^* = \infty$, $\infty^* = 0$. For fix $a \in \mathbb{B}^n \setminus \{0\}$, let

$$\sigma_a(z) = a^* + r^2(x - a^*)^*,\ r^2 = |a|^{-2} - 1 \tag{2.1}$$

be an inversion in the sphere $S^{n-1}(a^*, r)$ orthogonal to $S^{n-1}$ (see Figure 1). Then $\sigma_a(a) = 0$, $\sigma_a(a^*) = \infty$ and

$$|\sigma_a(x) - \sigma_a(y)| = \frac{r^2|x-y|}{|x-a^*||y-a^*|}. \tag{2.2}$$



FIGURE 1.

**Lipschitz mappings.** Let $(X, d_X)$ and $(Y, d_Y)$ be metric spaces. Let $f : X \to Y$ be continuous and let $L \geq 1$. We say that $f$ is $L$-lipschitz, if

$$d_Y(f(x), f(y)) \leq L d_X(x,y), \text{ for } x,\ y \in X,$$

and $L$-bilipschitz, if $f$ is a homeomorphism and

$$d_X(x,y)/L \leq d_Y(f(x), f(y)) \leq L d_X(x,y), \text{ for } x,\ y \in X.$$

A 1-bilipschitz mapping is called an isometry.

The so-called *monotone form of l'Hôpital's rule* is useful in deriving monotonicity properties and obtaining inequalities.

**Lemma 2.3.** [AVV, Theorem 1.25]. *For $-\infty < a < b < \infty$, let $f,\ g : [a,b] \to \mathbb{R}$ be continuous on $[a,b]$, and be differentiable on $(a,b)$, and let $g'(x) \neq 0$ on $(a,b)$. If $f'(x)/g'(x)$ is increasing(deceasing) on $(a,b)$, then so are*

$$\frac{f(x)-f(a)}{g(x)-g(a)} \quad \text{and} \quad \frac{f(x)-f(b)}{g(x)-g(b)}.$$

*If $f'(x)/g'(x)$ is strictly monotone, then the monotonicity in the conclusion is also strict.*

**Theorem 2.4.** *(1) Let $f : \mathbb{H}^2 \to \mathbb{B}^2$ with $f(z) = \frac{z-i}{z+i}$. Then for all $x, y \in \mathbb{H}^2$,*

$$j_{\mathbb{B}^2}(f(x), f(y)) \leq 2 j_{\mathbb{H}^2}(x,y),$$

*and the constant 2 is best possible.*

*(2) Let $f : \mathbb{H}^2 \setminus \{i\} \to \mathbb{B}^2 \setminus \{0\}$ with $f(i) = 0$, $f$ is the same as (1). Then for all $x, y \in \mathbb{H}^2 \setminus \{i\}$,*

$$j_{\mathbb{B}^2 \setminus \{0\}}(f(x), f(y)) \leq 2 j_{\mathbb{H}^2 \setminus \{i\}}(x,y),$$

*and the constant 2 is best possible.*

*Proof.* It suffices to show the sharpness of the inequalities by Theorem 1.1.

(1) Let $x = t + ai$ and $y = ai$, where $a > 0$ and $t > 0$. Then

$$\begin{aligned}
\lim_{t\to\infty} \frac{j_{\mathbb{B}^2}(f(t+ai), f(ai))}{j_{\mathbb{H}^2}(t+ai, ai)} &= \lim_{t\to\infty} \frac{\log(1 + \frac{t}{2a(a+1)}(\sqrt{t^2+(a+1)^2} + \sqrt{t^2+(a-1)^2})}{\log(1+\frac{t}{a})} \\
&= \lim_{t\to\infty} \frac{\log(\frac{2t^2}{2a(a+1)})}{\log(\frac{t}{a})} \\
&= 2.
\end{aligned}$$

Hence the constant 2 is best possible.

(2) Let $x = t + ai$ and $y = ai$, where $0 < a < \frac{1}{3}$ and $t > 1$. Then

$$\frac{j_{\mathbb{B}^2\setminus\{0\}}(f(x), f(y))}{j_{\mathbb{H}^2\setminus\{i\}}(x, y)} = \frac{j_{\mathbb{B}^2}(f(x), f(y))}{j_{\mathbb{H}^2}(x, y)}.$$

By (1) we obtain that the constant 2 is best possible. ☐

**Theorem 2.5.** *(1) Let* $f : \mathbb{B}^2 \to \mathbb{H}^2$ *with* $f(z) = i\frac{1+z}{1-z}$. *Then for all* $x$, $y \in \mathbb{B}^2$

$$j_{\mathbb{H}^2}(f(x), f(y)) \le 2 j_{\mathbb{B}^2}(x, y)\,,$$

*and the constant* 2 *is best possible.*

*(2) Let* $f : \mathbb{B}^2 \setminus \{0\} \to \mathbb{H}^2 \setminus \{i\}$ *with* $f(0) = i$, $f$ *is the same as (1). Then for all* $x$, $y \in G$

$$j_{\mathbb{H}^2\setminus\{i\}}(f(x), f(y)) \le 2 j_{\mathbb{B}^2\setminus\{0\}}(x, y)\,,$$

*and the constant 2 is best possible.*

*Proof.* By Theorem 1.1, we only need to show that 2 is best possible..

(1) For $t = x = -y,\ t \in (0, 1)$, we get

$$\begin{aligned}
j_{\mathbb{H}^2}(f(x), f(y)) &= \log(1 + \frac{4t}{(1-t)^2}) \\
&= 2\log(1 + \frac{2t}{1-t}) \\
&= 2 j_{\mathbb{B}^2}(x, y).
\end{aligned}$$

Therefore, the constant 2 is best possible.

(2) Since for $t = x = -y$ and $\frac{1}{2} < t < 1$ we have

$$j_{\mathbb{H}^2\setminus\{i\}}(f(x), f(y)) = j_{\mathbb{H}^2}(f(x), f(y)) = 2 j_{\mathbb{B}^2}(x, y) = 2 j_{\mathbb{B}^2\setminus\{0\}}(x, y),$$

the constant 2 is best possible. ☐

**Lemma 2.6.** *Let* $k \in (1, \infty)$, $\theta \in (0, \frac{\pi}{2k})$ *and* $r \in (0, 1)$. *Then*
*(1)* $f_1(\theta) \equiv \frac{k\sin\theta}{1-\sin\theta} - \frac{\sin k\theta}{1-\sin k\theta}$ *is decreasing from* $(0, \frac{\pi}{2k})$ *to* $(-\infty, 0)$.
*(2)* $f_2(r) \equiv \frac{1-(1-\sin\theta)r}{1-(1-\sin k\theta)r^k}$ *is decreasing from* $(0, 1)$ *to* $(\frac{\sin\theta}{\sin k\theta}, 1)$.
*(3)* $f_3(r) \equiv \frac{\log(1+\frac{1-r^k}{r^k \sin k\theta})}{\log(1+\frac{1-r}{r\sin\theta})}$ *is decreasing from* $(0, 1)$ *to* $(\frac{k\sin\theta}{\sin k\theta}, k)$.

*Proof.* (1) By differentiation,

$$f_1'(\theta) = k[g(1) - g(k)],$$

where $g(k) = \frac{\cos k\theta}{(1-\sin k\theta)^2}$. Since

$$g'(k) = \frac{\theta(1 - \sin k\theta + \cos^2 k\theta)}{(1 - \sin k\theta)^3} > 0,$$

we have $f_1'(\theta) < 0$ and hence $f_1$ is decreasing in $(0, \frac{\pi}{2k})$. The limiting values are clear.

(2) By differentiation,

$$\begin{aligned} & [1 - (1 - \sin k\theta)r^k]^2 f_2'(r) \\ = \; & -(k-1)(1 - \sin\theta)(1 - \sin k\theta)r^k + k(1 - \sin k\theta)r^{k-1} - (1 - \sin\theta) \\ \equiv \; & h(r). \end{aligned}$$

Since

$$h'(r) = k(k-1)(1 - \sin k\theta)r^{k-2}[1 - r(1 - \sin\theta)] > 0,$$

we have $h$ is increasing in $(0, 1)$ and hence $h(r) \leq h(1)$. By (1),

$$h(1) = (1 - \sin\theta)(1 - \sin k\theta)f_1(\theta) < 0.$$

Therefore, $f_2'(r) < 0$ and hence $f_2$ is decreasing in $(0, 1)$. The limiting values are clear.

(3) Let $f_3(r) \equiv \frac{g_1(r)}{g_2(r)}$, here $g_1(r) = \log(1 + \frac{1-r^k}{r^k \sin k\theta})$ and $g_2(r) = \log(1 + \frac{1-r}{r \sin\theta})$. Then $g_1(1^-) = g_2(1^-) = 0$, by differentiation, we have

$$\frac{g_1'(r)}{g_2'(r)} = kf_2(r).$$

Therefore, $f_3$ is increasing in $(0, 1)$ by Lemma 2.3. By l'Hôpital's Rule and (2) we get the limiting values easily. ☐

The planar angular domain is defined as

$$S_\varphi = \{re^{i\theta} \in \mathbb{C} : 0 < \theta < \varphi\,,\ r > 0\}.$$

**Lemma 2.7.** *Let* $f : S_{\pi/k} \to \mathbb{H}^2$ *with* $f(z) = z^k$ $(k \geq 1)$. *Let* $x, y \in S_{\pi/k}$ *with* $\arg(x) = \arg(y) = \theta$. *Then*

$$\frac{k \sin\theta}{\sin k\theta} j_{S_{\pi/k}}(x, y) \leq j_{\mathbb{H}^2}(f(x), f(y)) \leq k j_{S_{\pi/k}}(x, y).$$

*Proof.* Since $j-$metric is invariant under stretchings, we can assume that $x = re^{i\theta}$ and $y = e^{i\theta}$, $0 < r < 1$. By symmetry, we also assume $0 < \theta \leq \frac{\pi}{2k}$. Then

$$\frac{j_{\mathbb{H}^2}(f(x), f(y))}{j_{S_{\pi/k}}(x, y)} = \frac{\log(1 + \frac{1-r^k}{r^k \sin k\theta})}{\log(1 + \frac{1-r}{r \sin\theta})}.$$

By Lemma 2.6(3), we get the result. ☐

**Lemma 2.8.** *Let* $n \in \mathbb{N}$, $0 < \theta \leq \frac{\pi}{2n}$. *Then for* $x, y \in R^n \setminus \{0\}$,

$$1 + \frac{|x^n - y^n|}{|x|^n \sin n\theta} \leq (1 + \frac{|x - y|}{|x| \sin\theta})^n.$$

*Proof.* It is clear that (2.8) holds if $n = 1$.

Next, suppose that (2.8) holds when $n = k$ . Namely,

$$1 + \frac{|x^k - y^k|}{|x|^k \sin k\theta} \le (1 + \frac{|x-y|}{|x| \sin\theta})^k, \tag{2.9}$$

where $0 < \theta \le \frac{\pi}{2k}$.

Then, if $n = k+1$, we have $0 < \theta \le \frac{\pi}{2(k+1)} < \frac{\pi}{2k}$ and by (2.9),

$$\begin{aligned} (1 + \frac{|x-y|}{|x| \sin\theta})^{k+1} &\ge (1 + \frac{|x^k - y^k|}{|x|^k \sin k\theta})(1 + \frac{|x-y|}{|x| \sin\theta}) \\ &\ge 1 + \frac{|x-y|}{|x| \sin\theta} + \frac{|x^k - y^k|}{|x|^k \sin k\theta}(1 + \frac{|x-y|}{|x|}) \\ &\ge 1 + \frac{|x^{k+1} - x^k y|}{|x|^{k+1} \sin\theta} + \frac{|x^k y - y^{k+1}|}{|x|^{k+1} \sin k\theta} \\ &\ge 1 + \frac{|x^{k+1} - y^{k+1}|}{|x|^{k+1} \sin(k+1)\theta}. \end{aligned}$$

This completes the proof by induction. ☐

**Theorem 2.10.** *Let $f : S_{\pi/k} \to \mathbb{H}^2$ with $f(z) = z^k$ ($k \in \mathbb{N}$). Then for all $x, y \in S_{\pi/k}$,*

$$j_{\mathbb{H}^2}(f(x), f(y)) \le k j_{S_{\pi/k}}(x, y),$$

*and the constant $k$ is best possible.*

*Proof.* By symmetry, we can assume that $d(f(x), \partial\mathbb{H}^2) \le d(f(y), \partial\mathbb{H}^2)$. Hence by Lemma 2.8, we obtain

$$\begin{aligned} j_{\mathbb{H}^2}(f(x), f(y)) &= \log(1 + \frac{|x^k - y^k|}{|x|^k \sin k\theta}) \\ &\le k \log(1 + \frac{|x-y|}{|x| \sin\theta}) \\ &\le k j_{S_{\pi/k}}(x, y), \end{aligned}$$

where $0 < \theta = \min\{\arg(x), \frac{\pi}{k} - \arg(x)\} \le \frac{\pi}{2k}$.

Let $x = re^{i\alpha}$ and $y = e^{i\alpha}$, where $0 < \alpha < \frac{\pi}{2k}$ and $0 < r < 1$. Let $r \to 0$, then by Lemma 2.6(3) and Lemma 2.7, we know that the constant $k$ is best possible. ☐

**Lemma 2.11.** *For $z \in \mathbb{C}$ and $p \in \mathbb{N}$ we have*

$$\log(1 + |z^p - 1|) \le p \log(1 + |z - 1|).$$

*Proof.* Putting $z = u + 1$, we obtain

$$1 + |z^p - 1| = 1 + |\sum_{k=1}^{p} \binom{p}{k} u^k| \le 1 + \sum_{k=1}^{p} \binom{p}{k} |u|^k = (1 + |u|)^p = (1 + |z-1|)^p,$$

and the claim follows. ☐

**Lemma 2.12.** *Let $Q_d^*$ denote the class of all polynomials with exact degree $d$( $d \ge 1$) which have no zeros inside the disk $\mathbb{B}^2$.*

*If $Q \in Q_d^*$ and $u, v \in \mathbb{B}^2$, then*

$$|\frac{Q(u)}{Q(v)} - 1| \le \Big(1 + \frac{|u-v|}{1-|v|}\Big)^d - 1.$$

*Proof.* A polynomial $Q(z)$ with zeros $\{-a_k\}(k=1,2,\cdots,d)$ has a representation in the form

$$Q(z)=C\prod_{k=1}^{d}(z+a_k), C\neq 0.$$

Since $Q(z)\neq 0$ for $z\in\mathbb{B}^2$ then necessarily $|a_k|\geq 1$.

Therefore,

$$\begin{aligned}|\frac{Q(u)}{Q(v)}-1| &= |\prod_{k=1}^{d}\frac{u+a_k}{v+a_k}-1|\\ &= |\prod_{k=1}^{d}(1+\frac{u-v}{a_k+v})-1|\\ &= |\sum\prod(\cdot)(\cdot)|\leq\sum\prod|(\cdot)||(\cdot)|\\ &= \prod_{k=1}^{d}(1+\frac{|u-v|}{|a_k+v|})-1\\ &\leq \prod_{k=1}^{d}(1+\frac{|u-v|}{|a_k|-|v|})-1\\ &\leq \Big(1+\frac{|u-v|}{1-|v|}\Big)^d-1.\end{aligned}$$

$\square$

**Remark 2.13.** *Note that, since* $|\frac{Q(u)}{Q(v)}-1|\geq|\frac{Q(u)}{Q(v)}|-1$, *we also obtain*

$$|\frac{Q(u)}{Q(v)}|\leq\Big(1+\frac{|u-v|}{1-|v|}\Big)^d.$$

**Theorem 2.14.** *Let* $p\in\mathbb{N}$ *and* $\{a_k\}$ *be a sequence of complex numbers with* $\sum_{k=1}^{p}|a_k|\leq 1$. *Let* $f:\mathbb{B}^2\backslash\{0\}\to\mathbb{B}^2\backslash\{0\}$ *with* $f(z)=\sum_{k=1}^{p}a_kz^k$ *and* $f(0)=0$. *Then for all* $x,\ y\in\mathbb{B}^2\backslash\{0\}$

$$j_{\mathbb{B}^2\backslash\{0\}}(f(x),f(y))\leq pj_{\mathbb{B}^2\backslash\{0\}}(x,y),$$

*and the constant p is sharp.*

*Proof.* For $x,\ y\in\mathbb{B}^2\setminus\{0\}$ , we have

$$j_{\mathbb{B}^2\backslash\{0\}}(x,y)=\log(1+\frac{|x-y|}{\min\{|x|,|y|,1-|x|,1-|y|\}})$$

and

$$j_{\mathbb{B}^2\backslash\{0\}}(f(x),f(y))=\log(1+\frac{|f(x)-f(y)|}{T}),$$

where $T=\min\{|f(x)|,|f(y)|,1-|f(x))|,1-|f(y)|\}$.

Case 1. $T=1-|f(x)|$.

Since

$$\begin{aligned}|f(x)-f(y)| &= |x-y||\sum_{k=1}^{p}a_k(\sum_{i+j=k-1}x^iy^j)|\\ &\leq |x-y|\sum_{k=1}^{\infty}|a_k|(\sum_{i=0}^{k-1}|x|^i)\end{aligned}$$

and

$$\begin{aligned} 1-|f(x)| &\geq \sum_{k=1}^{p}|a_k| - \sum_{k=1}^{p}|a_k||x|^k \\ &= (1-|x|)\sum_{k=1}^{p}|a_k|(\sum_{i=0}^{k-1}|x|^i), \end{aligned}$$

we obtain

$$\begin{aligned} j_{\mathbb{B}^2\setminus\{0\}}(f(x),f(y)) &\leq \log(1+\frac{|x-y|}{1-|x|}) \\ &\leq j_{\mathbb{B}^2\setminus\{0\}}(x,y). \end{aligned}$$

Case 2. $T = 1-|f(y)|$.

It can be treated similarly to Case 1.

Case 3. $T = |f(y)|$.

Now if 0 is $m$-th order zero of $f$ and since $f$ obviously has no other zeros in $\mathbb{B}^2$, we get $f(z) = z^m Q(z),\ \ Q \in Q^*_d,\ m+d = p$, here $Q^*_d$ is as in Lemma 2.12.

By Lemma 2.12 and Remark 2.13, it follows

$$\begin{aligned} \frac{|f(x)-f(y)|}{|f(y)|} &= |\frac{x^m Q(x)}{y^m Q(y)} - 1| \\ &= |\Big(\frac{x^m}{y^m}-1\Big)\frac{Q(x)}{Q(y)} + \frac{Q(x)}{Q(y)} - 1| \\ &\leq |\frac{x^m}{y^m}-1||\frac{Q(x)}{Q(y)}| + |\frac{Q(x)}{Q(y)}-1| \\ &\leq \Big(1+|\frac{x^m}{y^m}-1|\Big)\Big(1+\frac{|x-y|}{1-|y|}\Big)^d - 1. \end{aligned}$$

Therefore, by Lemma 2.11 we have

$$\begin{aligned} j_{\mathbb{B}^2\setminus\{0\}}(f(x),f(y)) &= \log\Big(1+\frac{|f(x)-f(y)|}{|f(y)|}\Big) \\ &\leq d\log\Big(1+\frac{|x-y|}{1-|y|}\Big) + \log\Big(1+|\frac{x^m}{y^m}-1|\Big) \\ &\leq d\log\Big(1+\frac{|x-y|}{1-|y|}\Big) + m\log(1+\frac{|x-y|}{|y|}) \\ &\leq p j_{\mathbb{B}^2\setminus\{0\}}(x,y), \end{aligned}$$

and the proof for case 3 is done.

Case 4. $T = |f(x)|$.

It can be treated analogously to Case 3.

For the sharpness of the inequality, let $f(z) = z^p$ ( $p \in \mathbb{N}$). For $s$, $t \in (0, \frac{1}{2})$ and $s < t$, we have

$$j_{\mathbb{B}^2\setminus\{0\}}(f(t),f(s)) = \log(1+\frac{t^p - s^p}{s^p}) = p\log(\frac{t}{s}) = p j_{\mathbb{B}^2\setminus\{0\}}(t,s).$$

Therefore the constant $p$ is sharp. □

**Theorem 2.15.** *Let* $f(z) = a + r^2\frac{(z-a)}{|z-a|^2}$ *be the inversion in* $S^{n-1}(a,r)$ *with* $\operatorname{Im} a = 0$. *Then* $f(\mathbb{H}^n) = \mathbb{H}^n$ *and for* $x, y \in \mathbb{H}^n$,

$$j_{\mathbb{H}^n}(f(x), f(y)) \le 2 j_{\mathbb{H}^n}(x,y).$$

*The constant 2 is best possible.*

*Proof.* The inequality is clear by Theorem 1.1.

Putting $x = i$, $y = t+i$, $t > 0$, we have

$$\lim_{t\to\infty} \frac{j_{H^n}(f(i), f(t+i))}{j_{H^n}(i, t+i)} = \lim_{t\to\infty} \frac{\log(1+t\sqrt{1+t^2})}{\log(1+t)} = \lim_{t\to\infty} \frac{\log t^2}{\log t} = 2.$$

Hence the constant 2 is best possible.

□

## 3. PROOF OF MAIN RESULTS

In order to prove Theorem 1.4, we first need some lemmas.

**Lemma 3.1.** [B, Theorem 3.5.1, p.40]. *Let* $f$ *be a Möbius transformation and* $f(B^n) = B^n$. *Then*

$$f(x) = (\sigma x)A,$$

*where* $\sigma$ *is a inversion in some sphere orthogonal to* $S^{n-1}$ *and* $A$ *is an orthogonal matrix.*

**Lemma 3.2.** *Let* $a, b \in \mathbb{B}^n$. *Then*

*(1)*

$$|a|^2|b-a^*|^2 - |b-a|^2 = (1-|a|^2)(1-|b|^2).$$

*(2)*

$$\frac{||b|-|a||}{1-|a||b|} \le \frac{|b-a|}{|a||b-a^*|} \le \frac{|b|+|a|}{1+|a||b|}.$$

*Proof.* (1) By calculation, we have

$$\begin{aligned} & |a|^2|b-a^*|^2 - |b-a|^2 \\ = \ & |a|^2[|b|^2 + \frac{1}{|a|^2} - \frac{2(b\cdot a)}{|a|^2}] - [|b|^2 + |a|^2 - 2(b\cdot a)] \\ = \ & 1 + |a|^2|b|^2 - |a|^2 - |b|^2 \\ = \ & (1-|a|^2)(1-|b|^2). \end{aligned}$$

(2) This can be obtained by [Vu2, Exercise 2.52, p.32] directly. □

**Lemma 3.3.** *Let* $c, d \in (0,1), \theta \in (0,1]$. *Then*

*(1)* $f(\theta) \equiv \frac{\log\left(1+\frac{2cd\theta}{1-cd}\right)}{\log\left(1+\frac{2d\theta}{1-d}\right)}$ *is increasing. In particular,*

$$\frac{\log\left(1+\frac{2cd\theta}{1-cd}\right)}{\log\left(1+\frac{2d\theta}{1-d}\right)} \le \frac{\log\left(1+\frac{2cd}{1-cd}\right)}{\log\left(1+\frac{2d}{1-d}\right)}.$$

*(2)* $g(\theta) \equiv \frac{\operatorname{arth}(c\theta)}{\operatorname{arth}\theta}$ *is decreasing. In particular,*

$$\frac{\operatorname{arth}(c\theta)}{\operatorname{arth}\theta} \le c.$$

*(3)*

$$\Big(1+\frac{2cd\theta}{1-cd}\Big)\Big(1+\frac{c(1-d)}{1+cd}\Big)\le 1+\frac{c(1-d)+2cd\theta}{1-cd}\,.$$

*Proof.* (1) Let $f_1(\theta)=\log\Big(1+\frac{2cd\theta}{1-cd}\Big)$ and $f_2(\theta)=\log\Big(1+\frac{2d\theta}{1-d}\Big)$. Then we have $f_1(0^+)=f_2(0^+)=0$ and

$$\frac{f_1'(\theta)}{f_2'(\theta)}=1-\frac{1-c}{1-cd+2cd\theta},$$

which is clearly increasing in $\theta$. Therefore, the monotonicity of $f$ follows from Lemma 2.3. The inequality follows by the monotonicity of $f$ immediately.

(2) Let $g_1(\theta)=\operatorname{arth}(c\theta)$ and $g_2(\theta)=\operatorname{arth}\theta$. Then we have $g_1(0^+)=g_2(0^+)=0$ and

$$\frac{g_1'(\theta)}{g_2'(\theta)}=\frac{1}{c}(1-\frac{1-c^2}{1-c^2\theta^2})$$

which is clearly decreasing in $\theta$. Therefore, the monotonicity of $g$ follows from Lemma 2.3. The inequality follows by the monotonicity of $g$ and l'Hôpital's Rule immediately.

(3) It is easy to prove by direct calculation. ☐

Now we are in a position to give a short proof of Theorem 1.4.

**3.4. Proof of Theorem 1.4.** It is trivial for $a=0$, therefore, we only need to consider $a\neq 0$. Since $j$-metric is invariant under orthogonal transformations and by Lemma 3.1, for $x,y,a\in\mathbb{B}^n$, we have

$$j_{\mathbb{B}^n}(f(x),f(y))=j_{\mathbb{B}^n}(\sigma_a(x),\sigma_a(y)),$$

where $\sigma_a(x)$ is an inversion in the sphere $S^{n-1}(a^*,\sqrt{|a|^{-2}-1})$ orthogonal to $S^{n-1}$. Thus, it suffices to estimate the expression

$$J(x,y;a)\equiv\frac{j_{\mathbb{B}^n}(\sigma_a(x),\sigma_a(y))}{j_{\mathbb{B}^n}(x,y)}=\frac{\log\Big(1+\frac{|\sigma_a(x)-\sigma_a(y)|}{\min\{1-|\sigma_a(x)|,1-|\sigma_a(y)|\}}\Big)}{\log\Big(1+\frac{|x-y|}{\min\{1-|x|,1-|y|\}}\Big)}.$$

Let $r=\max\{|x|,|y|\}$ and suppose $|\sigma_a(x)|\ge|\sigma_a(y)|$. Then by (2.2), we have

$$\min\{1-|\sigma_a(x)|,1-|\sigma_a(y)|\}=1-|\sigma_a(x)|=\frac{|a||x-a^*|-|x-a|}{|a||x-a^*|}.$$

We first prove the right-hand side of inequality. By Lemma 3.2, we get

$$\begin{aligned}
j_{\mathbb{B}^n}(\sigma_a(x),\sigma_a(y)) &= \log\Big(1+\frac{(1-|a|^2)|x-y|}{|a||y-a^*|(|a||x-a^*|-|x-a|)}\Big)\\
&= \log\Big(1+\frac{|x-y|(|a||x-a^*|+|x-a|)}{|a||y-a^*|(1-|x|^2)}\Big)\\
&= \log\Big(1+\frac{|x-y||x-a^*|)}{(1-|x|^2)|y-a^*|}(1+\frac{|x-a|}{|a||x-a^*|})\Big)\\
&\le \log\Big(1+\frac{|x-y|}{1-r^2}(1+\frac{|x-y|}{|y-a^*|})(1+\frac{|x|+|a|}{1+|a||x|})\\
&\le \log\Big(1+\frac{|x-y|}{1-r}(1+\frac{|a||x-y|}{1-|a|r})(1+\frac{|a|(1-r)}{1+|a|r})\Big).
\end{aligned}$$

Then

$$\begin{aligned} J(x,y;a) &\leq \frac{\log\Big(1+\frac{|x-y|}{1-r}(1+\frac{|a||x-y|}{1-|a|r})(1+\frac{|a|(1-r)}{1+|a|r})\Big)}{\log\Big(1+\frac{|x-y|}{1-r}\Big)} \\ &= \frac{\log\Big(1+\frac{2r\theta}{1-r}(1+\frac{2|a|r\theta}{1-|a|r})(1+\frac{|a|(1-r)}{1+|a|r})\Big)}{\log\Big(1+\frac{2r\theta}{1-r}\Big)}, \end{aligned}$$

where $\theta = \frac{|x-y|}{2r}$.

By Lemma 3.3 it follows

$$\begin{aligned} J(x,y;a) &\leq \frac{\log(1+\frac{2r\theta}{1-r}(1+\frac{|a|(1-r)}{1-|a|r}+\frac{2|a|r\theta}{1-|a|r}))}{\log\Big(1+\frac{2r\theta}{1-r}\Big)} \\ &= 1+\frac{\log(1+\frac{2|a|r\theta}{1-|a|r})}{\log\Big(1+\frac{2r\theta}{1-r}\Big)} \\ &\leq 1+\frac{\log(1+\frac{2|a|r}{1-|a|r})}{\log\Big(1+\frac{2r}{1-r}\Big)} \\ &= 1+\frac{\text{arth}(|a|r)}{\text{arth}r} \\ &\leq 1+|a|. \end{aligned}$$

Therefore, we get

$$j_{\mathbb{B}^n}(f(x),f(y)) \leq (1+|a|)j_{\mathbb{B}^n}(x,y).$$

The sharpness of the upper bound $1+|a|$ is already proved in [KVZ].

For the left-hand side of inequality, we note that $f^{-1}(x) = A^{-1}\sigma_a^{-1}(x) = A^{-1}\sigma_a(x)$, here $\sigma_a(x)$ and $A$ are as above. Since $A$ is an orthogonal matrix and so is $A^{-1}$. Then by the above proof, for $x$, $y \in \mathbb{B}^n$, we get

$$\frac{j_{\mathbb{B}^n}(f^{-1}(x),f^{-1}(y))}{j_{\mathbb{B}^n}(x,y)} = \frac{j_{\mathbb{B}^n}(\sigma_a(x),\sigma_a(y))}{j_{\mathbb{B}^n}(x,y)} \leq 1+|a|.$$

Therefore, we have

$$j_{\mathbb{B}^n}(f(x),f(y)) \geq \frac{1}{1+|a|}j_{\mathbb{B}^n}(x,y). \tag{3.5}$$

This completes the proof. □

**3.6. Proof of Theorem 1.5.** A Möbius transformation satisfying the condition in Theorem 1.5 is of the form

$$f(z) = e^{i\alpha}\frac{z-a}{z-\bar{a}}$$

and hence

$$f^{-1}(z) = \frac{a-\bar{a}e^{-i\alpha}z}{1-e^{-i\alpha}z},$$

where $\alpha$ is a real constant. Since $j$-metric is invariant under translations, streching of $\mathbb{H}^2$ onto itself and rotations of $\mathbb{B}^2$ onto itself, we can assume that $a = i$ and $\alpha = 0$. Then we have

$$f(z) = \frac{z-i}{z+i} \quad \textit{and} \quad f^{-1}(z) = i\frac{1+z}{1-z}.$$

By Theorem 2.4 and Theorem 2.5, we get the results immediately. $\square$

**3.7. Proof of Theorem 1.6** Let $r = \max\{|x|, |y|\}$ and suppose that $|f(x)| \geq |f(y)|$. Then

$$j_{\mathbb{B}^2}(x,y) = \log(1 + \frac{|x-y|}{1-r})$$

and

$$j_{\mathbb{B}^2}(f(x), f(y)) = \log(1 + \frac{|f(x)-f(y)|}{1-|f(x)|}).$$

Similarly to the proof of Theorem 2.14 Case 1, we have

$$\begin{aligned} |f(x) - f(y)| &= |x-y| |\sum_{k=1}^{\infty} a_k (\sum_{i+j=k-1} x^i y^j)| \\ &\leq |x-y| \sum_{k=1}^{\infty} |a_k| (\sum_{i=0}^{k-1} |x|^i) \end{aligned}$$

and

$$\begin{aligned} 1 - |f(x)| &\geq \sum_{k=1}^{\infty} |a_k| - \sum_{k=1}^{\infty} |a_k| |x|^k \\ &= (1-|x|) \sum_{k=1}^{\infty} |a_k| (\sum_{i=0}^{k-1} |x|^i). \end{aligned}$$

Therefore,

$$j_{\mathbb{B}^2}(f(x), f(y)) \leq \log(1 + \frac{|x-y|}{1-|x|}) \leq j_{\mathbb{B}^2}(x,y).$$

For the sharpness of the inequality let $a_p = 1, a_i = 0, i \neq p$, i.e., $f(z) = z^p$ ( $p \in \mathbb{N}$). For $s$, $t \in (0,1)$ and $s < t$, we have

$$\begin{aligned} \lim_{t \to 1^-} \frac{j_{\mathbb{B}^2}(f(t), f(s))}{j_{\mathbb{B}^2}(t,s)} &= \lim_{t \to 1^-} \frac{\log \frac{1-s^p}{1-t^p}}{\log \frac{1-s}{1-t}} \\ &= \lim_{t \to 1^-} \frac{\log \frac{1-s}{1-t} + \log \frac{1+s+\cdots+s^{p-1}}{1+t+\cdots+t^{p-1}}}{\log \frac{1-s}{1-t}} \\ &= 1. \end{aligned}$$

Therefore the inequality is sharp and the sharpness also clearly holds for identity. $\square$

**Conjecture 3.8.** *Let $a \in \mathbb{B}^2$ and $f : \mathbb{B}^2 \setminus \{0\} \to \mathbb{B}^2 \setminus \{a\}$ be a Möbius transformation with $f(0) = a$. Then for $x$, $y \in \mathbb{B}^2 \setminus \{0\}$*

$$j_{\mathbb{B}^2 \setminus \{a\}}(f(x), f(y)) \leq C(a) j_{\mathbb{B}^2 \setminus \{0\}}(x,y),$$

*where the constant $C(a) = 1 + (\log \frac{2+|a|}{2-|a|})/\log 3$ is best possible.*

**Acknowledgments.** The research of Matti Vuorinen was supported by the Academy of Finland, Project 2600066611. The research of Gendi Wang was supported by CIMO of Finland, Grant TM-10-7364.

## REFERENCES

[AVV] G.D. ANDERSON, M.K. VAMANAMURTHY, AND M. VUORINEN: *Conformal Invariants, Inequalities and Quasiconformal Maps*, John Wiley & Sons, New York, 1997.

[B] A. F. BEARDON: The geometry of discrete groups, Graduate Texts in Math. 91, Springer-Verlag, New York, 1995.

[GO] F. W. GEHRING AND B. G. OSGOOD: Uniform domains and the quasihyperbolic metric, J. Analyse Math. 36, 1979, 50-74.

[GP] F. W. GEHRING AND B.P. PALKA: Quasiconformally homogeneous domains, J. Analyse Math. 30, 1976, 172-199.

[HIMPS] P. HÄSTÖ, Z. IBRAGIMOV, D. MINDA, S. PONNUSAMY AND S. K. SAHOO: Isometries of some hyperbolic-type path metrics, and the hyperbolic medial axis, In the tradition of Ahlfors-Bers, IV, Contemporary Math. 432, 2007, 63-74.

[K1] R. KLÉN: Local convexity properties of quasihyperbolic balls in punctured space. - J. Math. Anal. Appl. 342:1, 2008, 192–201.

[K2] R. KLÉN: Local convexity properties of j-metric balls. - Ann. Acad. Sci. Fenn. Math. 33:1, 2008, 281–293.

[KVZ] R. KLÉN, M. VUORINEN, AND X. ZHANG: Quasihyperbolic metric and Möbius transformations.- Manuscript 9pp, August 2011. arXiv: 1108.2967 math.CV.

[MV] O. MARTIO AND J. VÄISÄLÄ: Quasihyperbolic geodesics in convex domains II. - Pure Appl. Math. Q. 7, 2011, 395–409.

[RT] A. RASILA AND J. TALPONEN: Convexity properties of quasihyperbolic balls in Banach spaces. Ann. Acad. Sci. Fenn. Math. 37, 2012, 215–228.

[V1] J. VÄISÄLÄ: The free quasiworld. Freely quasiconformal and related maps in Banach spaces. - In: Quasiconformal geometry and dynamics (Lublin, 1996), Banach Center Publ. 48, Polish Acad. Sci., Warsaw, 1999, 55–118.

[V2] J. VÄISÄLÄ: Quasihyperbolic geometry of planar domains. - Ann. Acad. Sci. Fenn. Math. 34, 2009, 447–473.

[Vu1] M. VUORINEN: *Conformal invariants and quasiregular mappings.* - J. Anal. Math. 45 (1985), 69-115.

[Vu2] M. VUORINEN: Conformal geometry and quasiregular mappings. Lecture Notes in Math. 1319, Springer-Verlag, Berlin-Heidelberg, 1988.

MATHEMATICAL INSTITUTE SANU, KNEZA MIHAILA 36, 11000 BELGRADE, SERBIA
*E-mail address*: `ssimic@turing.mi.sanu.ac.rs`

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TURKU, TURKU 20014, FINLAND
*E-mail address*: `vuorinen@utu.fi`
*E-mail address*: `genwan@utu.fi`